Utah Health Status by Race/Ethnicity, Summary Sheet

 ↑ =significantly higher than state overall =difference not statistically significant ↓ =significantly lower than state overall 	Utah, Overall	American Ind./ Alaska Native	Asian	Black	Native HI/ Pacific Islander	White	Hispanic	
Demographic Context								
Age Distribution of the Population, 2000	27.1	23.2	29.6	24.8	21.4	27.8	23.0	
* All Cause Death Rates, 1998-2003	797.2	872.0	352.3 [§] ↓	1,016.0 🛧	352.3§ ◆	796.3	710.3	
Life Expectancy at Birth, 1998-2003	77.9	74.5	84.9 [§]	74.2	84.9 [§]	78.0	78.2	
* Years of Potential Life Lost, 1998-2000	6,248	9,969	5,631 [§]	9,589	5,631 [§]	6144 [†]	6,326	
Poverty (All Ages), 2000	9.1%	17.4%	4.6%	8.2%	7.7%	8.7%	12.7%	
Child Poverty, 2000	10.1%	37.6%	13.1%	23.9%	18.3%	8.5%	22.2%	
Health Care Services and Systems								
* % Utahns With No Health Insurance, 2001	8.2%	12.1%	5.8%	7.3%	7.6%	7.1% 	24.8%	
* % of Insured Unable to Get Needed Health Care, 2001	12.2%	17.3%	8.7%	10.5%	4.6% ♥	12.2%	14.0%	
* % Utahns With No Usual Place of Medical Care, 2001	8.6%	10.9%	7.0%	12.0%	9.3%	7.7% ↓	20.6%	
* % Utahns Who Had a Routine Medical Check-up, 2001	71.0%	74.7%	76.5%	72.3%	78.9%	70.3%	80.2% 🛧	
* Average Number of Medical Visits, 2001	3.8	5.2	3.8	3.7	4.0	3.9	3.7	
% Age 50+, Colon Cancer Screening, 1999-2004	37.2%	37.8%	47.3%	57.4%	19.9%	37.2%	27.4% ↓	
* % Women, Pap Test, 1999-2000, 2002, 2004	80.7%	71.3%	85.6%	81.8%	71.1%	80.8%	72.7% •	
* %Women 40+, Mammogram, 1999-2000, 2002, 2004	67.7%	60.9%	78.9%	79.6%	55.5%	67.9%	61.7%	
* % Men 40+, PSA Test, 2001-2004	55.2%	43.1% ↓	58.8%	37.4%	69.4%	56.1%	42.1% ↓	
* % Adults Had Cholesterol Checked, 1999, 2001, 2003	67.6%	60.7%	70.3%	61.5%	63.3%	68.1%	58.7% ↓	
* % Adults With High Cholesterol, 1999, 2001, 2003	21.4%	13.6%	25.0%	17.2%	17.4%	21.6%	20.4%	
* % Adults Had Blood Pressure Checked, 1999	92.5%	83.8%	87.0%	100.0%	100.0%	92.7%	92.7%	
* % Adults With High Blood Pressure, 1999, 2001, 2003	22.6%	28.4%	24.6%	35.8%	28.6%	22.5%	22.8%	
% Mothers Had Early Prenatal Care, 2002	78.0%	53.9% ◆	72.5% 	57.2% ₩	48.1% ◆	79.1% ↑	60.4% ◆	
* % Adults Had Flu Shot, 1999, 2001-2004	35.2%	38.1%	45.0% ↑	32.1%	36.4%	35.1%	31.3% ↓	
Risk Factors For Illness								
* % Adults Overweight or Obese, 1999-2004	55.6%	66.3%	32.1% ↓	71.6%	79.9% ↑	55.2%	64.0%	
* % Adults Eating 2+ Fruits Daily, 1999-2000, 2002-2003	32.0%	29.2%	41.1%	34.3%	37.4%	31.7%	32.7%	
* % Adults Eating 3+ Veg. Daily, 1999-2000, 2002-2003	22.0%	19.1%	31.0%	15.8%	19.6%	22.1%	16.4% ♥	
* % Adults With No Physical Activity, 1999-2004	17.7%	33.4%	19.3%	25.2%	9.1% ↓	17.0% ₩	28.0%	





Utah Health Status by Race/Ethnicity, Summary Sheet (continued)

=significantly higher than state overall =difference not statistically significant =significantly lower than state overall	Utah, Overall	, l		Asian		Black		Native HI/ Pacific Islander		White		Hispan	ic
sk Factors For Illness (continued)													
* % Adults With Physical Activity, 2001, 2003	54.6%	54.6%		40.3%	4	58.6%		67.1%		55.5%		47.8%	Ψ
* % Adults Reported Cigarette Smoking, 1999-2004	12.3%	18.6%		8.4%		28.4%	1	6.7%	Ψ	12.0%		14.1%	
* % Adults Reported Chronic Drinking, 1999, 2001-04	3.3%	8.6%	↑	0.0%	4	3.7%		0.0%		3.3%		4.2%	
* % Adults Reported Binge Drinking, 1999, 2001-2004	9.4%	18.5%	↑	3.6%	Ψ	24.9%	1	3.9%		8.9%		14.5%	↑
* % Adults Reported DUI, 1999, 2002, 2004	0.9%	2.6%		0.0%		1.3%		0.0%		0.9%		1.4%	
* % Adults Not Always in Seat Belt	9.3%	11.2%		18.9%		0.0%		10.9%		9.1%		8.4%	
ealth Problems of Mothers and Infants													
Infant Mortality, 1998-2003	5.2	5.0		4.6 [§]		13.8	1	4.6 [§]		5.1		6.4	1
Infnat Mortality Related to Birth Defects, 1999-2003	1.5	1.3		1.4 [§]		1.3		1.4 [§]		1.5		1.4	
% Infants Low Birth Weight, 2002	6.6%	9.2%	↑	8.8%	↑	14.7%	1	7.1%		6.5%		6.6%	
Births to Adolescents (Age 15-17), 2002	17.5	44.0	↑	23.0		29.2		31.9	1	16.8		65.9	1
Overall Birth Defects, 1999-2003	20.6	17.7		17.2 [§]	Ψ	14.3	4	17.2 [§]	Ψ	21.2		18.4	¥
* % Women 18-44 Taking Folic Acid, 1999-2004	48.3%	36.6%		48.7%		34.9%		25.0%	Ψ	49.5%		37.7%	¥
Neural Tube Defects, 1994-2003	7.3	9.5		3.8 [§]	Ψ	0.0		3.8 [§]	Ψ	7.0		10.5	1
Orofacial Clefts, 1995-2003	22.2	36.9		18.2 [§]		12.0		18.2 [§]		22.2		22.4	
Congenital Heart Defects, 2003	62.9	66.8		71.6 [§]		0.0		71.6 [§]		61.5		74.4	
fectious Diseases	•												
Hepatitis A, 1998-2002	4.1	1.8		3.7 [§]		2.6		3.7 [§]		2.9	4	6.4	
Tuberculosis, 2000-2004	1.7	4.2		15.8	↑	19.9	1	10.3	↑	1.1	+	5.9	↑
Chlamydia, 2000-2004	147.5	212.3	↑	158.1 [§]		368.6	1	158.1 [§]		92.8	4	402.1	↑
Syphilis, 2000-2004	0.4	3.6	↑	1.3 [§]		1.7		1.3 [§]		0.2		0.7	
Gonorrhea, 2000-2004	16.5	18.4		12.8 [§]		85.0	1	12.8 [§]		11.4	4	34.6	↑
HIV/AIDS, 2000-2004	7.7	8.3		4.7 [§]		78.0	1	4.7 [§]		5.3	4	15.7	1
ury & Violence													
* % Utahns With Injury, 2001	11.9%	10.7%	•	8.2%	•	9.1%		13.2%		12.3%		9.1%	Ψ
* Unintentional Injury Deaths, 1998-2003	33.8	76.7	↑	15.3 [§]	Ψ	28.8		15.3 [§]	4	33.2		34.1	
* Motor Vehicle Traffic Crash Deaths, 1998-2003	15.4	52.4	↑	7.2 [§]	Ψ	17.6		7.2 [§]	4	14.7		18.6	

Utah Health Status by Race/Ethnicity, Summary Sheet (continued)

 ↑ =significantly higher than state overall =difference not statistically significant ↓ =significantly lower than state overall 	Utah, Overall	American Ind./ Alaska Native		Asian		Black		Native HI/ Pacific Islander		White	Hispanic	
Injury & Violence (continued)												
* Work-related Injury Deaths, 1998-2003	2.3	2.9		0.0§		1.5		0.0§		2.3	3.0	
* Suicide, 1998-2003	15.5	10.1	4	2.6 [§]	4	7.8	4	2.6 [§]	4	15.8	12.0	
* Homicide, 1998-2003	2.6	6.6	↑	0.7 [§]	4	9.4	1	0.7 [§]	4	2.4	7.4	↑
Chronic Diseases and Conditions												
* % Utahns in Fair/Poor Health (All Ages), 2001	11.0%	21.5%	↑	7.5%		13.2%		17.6%	↑	10.5%	19.5%	↑
* % Adults With Poor Physical Health, 1999-2004	14.4%	20.5%		9.5%		18.0%		10.6%		14.1%	16.4%	
* % Adults With Poor Mental Health, 1999-2004	15.0%	22.9%	↑	9.9%	4	27.1%	↑	15.9%		14.7%	16.9%	
* % Adults With Arthritis, 2000-2003	23.8%	32.2%		14.1%	4	34.3%		12.1%		23.9%	22.9%	
* % Utahns With Asthma (All Ages), 2001	5.5%	11.4%	↑	6.7%		6.3%		6.2%		5.7%	4.4%	
* % Utahns With Diabetes (All Ages), 2001	4.5%	8.0%	↑	5.2%		7.0%		5.8%		4.4%	5.6%	
* Diabetes Deaths, 1998-2003	73.0	147.4	↑	50.0 [§]		113.7		50.0 [§]		71.9	99.8	
* % Utahns With Coronary Heart Disease (All Ages), 2001	4.5%	8.7%	↑	5.4%		7.9%		5.7%		4.5%	3.9%	
* Coronary Heart Disease Deaths, 1998-2003	99.4	93.6		31.9 [§]	4	103.9		31.9 [§]	+	99.9	71.7	•
* Stroke Deaths, 1998-2003	56.2	22.2	4	23.0 [§]	4	83.1		23.0 [§]	+	56.2	49.9	
Cancer												
* Lung Cancer Incidence, 1997-2001	30.5	15.9		26.4		15.7		24.2		30.7	40.0	
* Lung Cancer Deaths, 1998-2003	24.8	19.1		19.5 [§]		36.0		19.5 [§]		24.8	24.2	
* Colorectal Cancer Incidence, 1997-2001	41.3	11.0	4	32.2		50.3		23.8		41.6	48.2	
* Colorectal Cancer Deaths, 1998-2003	16.1	7.9		10.8 [§]		35.8	↑	10.8 [§]		16.1	14.5	
* Breast Cancer Incidence (Females), 1997-2001	117.3	17.9	4	73.2		38.3	Ψ	36.5	+	119.6	111.5	
* Breast Cancer Deaths (Females), 1998-2003	23.1	15.0		6.2 [§]	Ψ	21.0		6.2 [§]	4	23.2	9.7	Ψ
* Prostate Cancer Incidence (Males), 1997-2001	176.7	54.3	4	62.5	Ψ	255.6		100.8		179.0	168.0	
* Prostate Cancer Deaths (Males), 1998-2003	31.4	17.0		11.5 [§]	Ψ	63.0		11.5 [§]	4	31.5	35.3	

^{*} Age adjusted to 2000 U.S. population



[§] Asian and Pacific Islander were grouped together.

[†] White, non-Hispanic